

Convergent Approximate Solving of First-Order Constraints by Approximate Quantifiers

STEFAN RATSCHAN

Research Institute for Symbolic Computation
Universität Linz

February 1, 2008

Abstract

Exactly solving first-order constraints (i.e., first-order formulas over a certain predefined structure) can be a very hard, or even undecidable problem. In continuous structures like the real numbers it is promising to compute approximate solutions instead of exact ones. However, the quantifiers of the first-order predicate language are an obstacle to allowing approximations to arbitrary small error bounds. In this paper we remove this obstacle by modifying the first-order language and replacing the classical quantifiers with approximate quantifiers. These also have two additional advantages: First, they are tunable, in the sense that they allow the user to decide on the trade-off between precision and efficiency. Second, they introduce additional expressivity into the first-order language by allowing reasoning over the size of solution sets.

1 Introduction

Solving first-order constraints, (i.e., first-order formulas over a certain predefined structure), and especially first-order constraints over the reals. has numerous applications [35, 13, 5, 12, 25, 28]. However, solving such constraints over the reals is either highly complex (e.g., when considering the predicate symbols $=$ and $\leq$, and the function symbols $+$ and $\times$ [39, 14, 11, 45]), or impossible [39, 37]. To deal with this problem, one can either restrict one-selves to more special problem classes (see e.g. [46, 23, 18]), or relax the problem by allowing approximation up to a user-specified error bound (as proposed by H. Hong [24]). This paper studies the general feasibility of the second approach. Its main contributions are: To show that even for this relaxed specification we might have to do exact intermediate computation; and to

introduce a modification of the first-order predicate language—approximate quantifiers—for which this problem does not occur.

These quantifiers have two additional advantages: First, they are tunable, in the sense that they allow the user to decide on the trade-off between precision and efficiency. Second, they introduce additional expressivity into the first-order predicate language, by allowing reasoning over the size of solution sets.

The first step to introduce approximate quantifiers is to allow quantifiers with a positive real annotation q , with the intuitive meaning that a formula $\exists_q x \phi$ is true iff the volume of the solution set of ϕ is greater than q . We will see that this does not yet allow the computation of approximate solution sets up to arbitrarily small, user-specified, error bounds.

So we allow quantifiers to be annotated with a real interval $[q, \bar{q}]$, with the intuitive meaning that the exact annotation can be any element of $[q, \bar{q}]$. This allows an algorithm to choose the most suitable value in $[q, \bar{q}]$ and we do not care which one. This means that a sentence in the language does not have one distinct truth-value but has a set of possible truth-values (cmp. with [20] or [22]). We will prove that, from a good enough approximation of the solution sets of the atomic sub-constraints, we can always compute at least one of these truth-values, and thus one can always attain an arbitrarily small error bound when computing approximate solution sets.

Following the usual approach (see e.g. [43]), one would implement such a logic using sets of truth-values (representing a many-valued logic [40, 36]) instead of single truth-values. We show that this approach is not suitable here and present a new method that is completely orthogonal to the semantics usually given to formulas when we do not know the value of certain predicate and function symbols and thus assign validity to the formula using all possible predicate and function symbol assignments (i.e., interpretations). Although arising from problems over real numbers, the resulting first-order language is completely domain-independent.

The above situation that there are several possible values for an object, and we either do not know or do not care which one should be taken, is commonly called *don't know* and *don't care nondeterminism*, respectively. Here both forms occur at the same time, which creates various difficulties through their interaction. Our approach gives general insight into such a situation by showing how one can compute with such nondeterministic objects, nevertheless. So our language can be easily extended to take into account nondeterminism coming from other sources. For example uncertain coefficients of occurring polynomials can be either modeled as don't care nondeterminism (the *united* approach) or as don't know nondeterminism (the *robust* approach) [8, 16, 38].

The structure of the paper is as follows: In Section 2, we give the specification of solving first-order constraints approximately up to some user-specified error bound, and informally show that this is impossible for con-

straints containing classical quantifiers. In Section 3, we develop a suitable formal model for approximate computation. In Section 4 we use this model to formalize the approximate solving of first-order constraints and its problems discussed in Section 2. In Section 5, we give a first-order language where the classical quantifiers are replaced by approximate ones. In Section 6, we apply the formal model for approximately solving first-order constraints to approximate quantifiers. In Section 7, we show how to deal with the resulting two forms of nondeterminism. In Section 8, we prove that one can solve constraints that contain approximate instead of classical quantifiers up to an arbitrary small error bound. In Section 9, we discuss related work, and in Section 10, we give a final conclusion.

2 Approximate Solving of First-Order Constraints

Throughout the paper we use the term “constraint” as a shortcut for “first-order constraint”, that is, a first-order formula over a certain, predefined structure $\mathcal{S}$. We fix a set V of variables and define a *variable assignment* as a function from V to $\mathcal{S}$. For a variable assignment θ , an element $a \in \mathcal{S}$, and variable $v \in V$, θ_v^a is the variable assignment that is the same as θ except that it assigns a to v .

We define the property that a constraint *is true* for a certain variable assignment (or *is satisfied by it*) as usual. A *potential solution set* is a set of variable assignments (we use the adjective “potential” for signifying the independence from a specific constraint), and the *solution set of a first-order constraint* ϕ is the set of variable assignments for which ϕ is true.

Sometimes we denote the solution set of a closed first-order constraint (i.e., *sentence*) by the Boolean constant $\mathbf{T}$ (which represents the set of all variable assignments), or $\mathbf{F}$ (which represents the empty set). In this case we also speak of the *truth-value* (instead of solution set) of a first-order constraint.

Recall that the notion of volume is modeled in mathematics by measure spaces (see any textbook on measure theory, for example [19], for details). For any measure μ and set A , the inner measure $\underline{\mu}(A)$ is the supremum of the measures of all measurable subsets of A (or $-\infty$, if this supremum does not exist), and the outer measure $\overline{\mu}(A)$ is the infimum of the measures of all measurable supersets of A (or ∞ , if this infimum does not exist). For any set A , $\underline{\mu}(A) \leq \overline{\mu}(A)$. If A is measurable, then $\mu(A) = \underline{\mu}(A) = \overline{\mu}(A)$ —in this case we often use the term *volume* for measure. Furthermore, we call a function μ *possible measure* iff for any set A , $\mu(A) \in [\underline{\mu}(A), \overline{\mu}(A)]$. We straightforwardly extend any measure on $\mathcal{S}^{|V|}$ to a measure on (potential) solution sets by measuring the tuples corresponding to the variable assignments in the (potential) solution set.

We want to solve constraints: Given some constraint, we want to get a

simple (e.g., quantifier-free) representation of its solution set. However, over the real numbers, this problem is either highly complex [39, 14, 11, 45] or undecidable [39, 37]. So we can only hope to tackle the general problem, if we relax it. As proposed by H. Hong [24], we do this by allowing approximation. This results in the problem specification of Figure 1.

GIVEN:	A constraint ϕ , and a positive real error bound ε
FIND:	Sets Y and N of variable assignments, such that ϕ is true for all elements of Y , ϕ is false for all elements of N , the volume of the variable assignments not in Y or N is smaller than ε

Figure 1: Problem Specification

In this paper we assume that we already have an algorithm that implements this specification for atomic input constraints. For example, one can use for this a branch-and-bound approach based on interval arithmetic [27]. In theory, one would need arbitrary precision here. In practice, however, fast machine-precision floating-point arithmetic usually suffices.

We would like to find an algorithm that fulfills the given specification based on such a solver for atomic constraints. However, this is impossible in general, because sometimes exact solution sets of atomic sub-constraints are needed to compute such approximate solutions. We show this here informally and formalize the arguments in the two following sections. Take an input constraint of the form $\exists x \phi$ without free variables, where ϕ is an atomic constraint with an empty solution set. Determine for all x , except for a set of arbitrarily small but positive volume, whether x is in the solution set of ϕ . From this information we cannot deduce that $\exists x \phi$ is false, because some of the remaining x might be in the solution set of ϕ , in which case $\exists x \phi$ would be true.

One could suspect that the reason for this problem is, that a classical quantifier has to take into account arbitrarily small solution sets of the quantified sub-constraint. So let us introduce an additional quantifier $\exists_q$ (the *existential volume quantifier*) into the first-order predicate language, where q is a non-negative real number. A closed constraint of the form $\exists_q x \phi$ is true iff the volume of the x , for which ϕ is true, is greater than q . But even for this quantifier, we have the same problem: Take an input constraint of the form $\exists_q x \phi$. Assume that the solution set of ϕ has exactly the volume q . Determine for all x , except for a set of arbitrarily small but

positive volume, whether x is in the solution set of ϕ . Again, from this information we cannot deduce whether $\exists_q x \phi$ is true. Note that one can easily find examples of constraints *with* free variables, that show the same behavior; in this case we can deduce for no element of the free-variable space whether it is an element of the solution set of such a constraint.

Speaking in the language of numerical analysis, the problem of finding approximations of solution sets of quantified constraints is *ill-posed* for certain inputs. This means that one can only solve it by approximation methods either after using more information, or after relaxing the problem. Following the latter approach, we introduce different quantifiers, for which the problem does not occur.

3 A Model for Approximate Computation

Before solving the problems described in the last section, we formalize them. For this we develop a model for approximate computation in this section, and apply it to approximate solving of first-order constraints in the next section. Readers who want to see a solution to the problem immediately, without being interested in formal proofs, can skip these sections and can directly jump to Section 5.

Assume a set A . Instead of doing exact computation in A , we use a set $\hat{A}$ for approximating computation in A . For this we add a notion of error, that is, a function from $\hat{A}$ to $\mathbb{R}^+$. For example, one can do approximate computation for the real numbers using the set of rational intervals. Here the error of an interval is its width. Take the expression $2x + 1$. If we know that x is in the interval $[2, 3]$ then we can deduce that the value of the expression is within $[5, 7]$ —a result with error 2.

Here we would like to be able to make the output error arbitrarily small by making the input error small:

Definition 1 *A function $f : \hat{A} \rightarrow \hat{A}$ is convergent iff for all $\varepsilon \in \mathbb{R}^+$ there is a $\delta \in \mathbb{R}^+$ such that for all $\hat{a} \in \hat{A}$ such that the error of $\hat{a}$ is less or equal δ , the error of $f(\hat{a})$ is less or equal ε .*

Note that this definition corresponds to the definition of uniform continuity in analysis; this notion and similar ones are used as necessary conditions for computability in effective analysis [6, 7, 33, 44].

We assume that every element $\hat{a} \in \hat{A}$ gives us the information that the result of some computation is in a certain subset of A . So we use subsets of A for modeling approximate computation from now on and simply identify $\hat{A}$ with 2^A . As in the example of rational intervals, very often one just uses certain subsets of A that allow a convenient representation.

Observe that it can also occur that we cannot define a certain function exactly, but only approximately. This means that there are several possible

functions, and we do not know which one is the correct one to use. Again we represent this *don't know* nondeterminism by a set of functions, which we call *approximate function*. For example, the function computing the weight of a certain mass depends on the constant describing gravitational acceleration. The exact value of this constant depends on the distance from the center of the earth, and so we have not exact value for it—only an interval covering all its possible values on the surface of the earth. Therefore several such functions are possible and the whole function is approximate.

If we want to do approximate computation, then the best possible result we can obtain without producing wrong results, is:

Definition 2 *For an approximate function $\hat{f}$ such that each element is a function on A , for an $\hat{a} \in \hat{A}$*

$$Ext_{\hat{f}}(\hat{a}) := \{f(a) \mid f \in \hat{f}, a \in \hat{a}\}$$

We call $Ext_{\hat{f}}$ the extension of $\hat{f}$.

For example, interval arithmetic [30, 32] defines approximate computation on the real numbers in this way. However, instead of computing with exact rational end-points, the results are usually rounded to the smallest super-interval whose end-points are floating-point numbers.

Clearly the fact that the extension of a function is not convergent, implies that we cannot use approximate computation to compute this function up to an arbitrarily small error.

4 Approximate Computation and Classical Quantifiers

Now we use the tools developed in the last section to show that classical quantifiers are an obstacle to approximate computation. Let $\mathbb{S}$ be the set of potential solution sets. Then we can do approximate computation using $\hat{\mathbb{S}}$, whose elements we call *potential approximate solution sets*, where the *approximate solution set of a constraint ϕ* is a potential approximate solution set that contains the solution set of ϕ .

We can deduce the information we want to find for our specification (Figure 1) as follows:

Definition 3 *A variable assignment θ is determined to true by a potential approximate solution set $\hat{d}$ iff for all $d \in \hat{d}$, $\theta \in d$. It is determined to false by $\hat{d}$ iff for all $d \in \hat{d}$, $\theta \notin d$. It is determined by $\hat{d}$ iff it is either determined to true or determined to false by $\hat{d}$.*

Now we can easily define the error of a potential approximate solution set $\hat{d}$ as the upper measure of the set of all variable assignments that are not determined by $\hat{d}$.

For example, an approximate solution set of the constraint $x^2 + y^2 \leq 1$ might contain all the potential solution sets that do not contain elements outside of the rectangle $[-1, 1] \times [-1, 1]$ (the solution set of $x^2 + y^2 \leq 1$ is one of these). In this case all variable assignments assigning values outside of $[-1, 1]$ are determined to false, and no variable assignment is determined to true. By measuring the size of the rectangle $[-1, 1] \times [-1, 1]$ we get the error 4 of this approximate solution set.

For a constraint of the form $L(\phi_1, \dots, \phi_n)$, where L is either a quantifier and a variable (in this case $n = 1$) or a connective, the solution set of the total constraint is a function of the solution sets of the sub-constraints $\phi_1, \dots, \phi_n$. It is easy to show that the functions corresponding to connectives are convergent in the sense of Definition 1. So we will concentrate on the case where L contains an (existential) quantifier. In this case we have the following function on potential solution sets (see [34] for the other cases).

Definition 4 *For a variable $v \in V$, the v -projection operator is a function P on potential solution sets such that a variable assignment $\theta \in P(d)$ iff there is an $a \in \mathcal{S}$ such that $\theta_v^a \in d$.*

So, for a constraint $\exists x \phi$, we can compute an approximate solution set of the total constraint from an approximate solution set of ϕ , by applying $\text{Ext}_{\{P\}}$, where P is the x -projection operator, to the approximate solution set of ϕ . Extending the above example to the constraint $\exists y x^2 + y^2 \leq 1$, we can use the y -projection operator to compute an approximate solution set of this constraint. The result contains all potential solution sets that do not assign values less than -1 or greater than 1 to x . This determines all variable assignments for which x is not in $[-1, 1]$ to false, and leaves all other variable assignments undetermined.

Using the argument from Section 2 it is easy to show that the extension of the approximate function that contains the projection operator is not convergent. Thus we cannot fulfill our specification (Figure 1) for constraints containing quantifiers. So we have to find a similar, but easier problem for which we can fulfill the specification. We do this by introducing quantifiers that approximate classical quantification, but result in a convergent projection operator.

5 A First-Order Predicate Language with Approximate Quantifiers

Classical quantifiers and volume quantifiers do not allow approximation up to an arbitrarily small error bound because they discontinuously change from

false to true. If the size of the solution set of the quantified constraint is exactly at this point of discontinuous change, then we cannot decide between true and false using approximate computation.

We avoid this discontinuous change by using volume quantifiers $\exists_q$ ($\forall_q$) for which we do not specify the annotation q exactly, but only require it to be within a certain interval $[\underline{q}, \bar{q}]$. Then a constraint solver can choose an element of this interval element for which it can safely decide whether the total constraint is true, using the available approximate information. This means that sentences containing such an approximate quantifier can possibly be both true and false—depending on which element of the interval $[\underline{q}, \bar{q}]$ we choose. This happens if the size of the solution set of a quantified constraint is within the interval $[\underline{q}, \bar{q}]$ (see Figure 2).




Figure 2: Approximate Quantifiers

Recall that an approximate solution set of a constraint consists of several potential solution sets, and we do not know which of them is the correct one—we have *don't know* nondeterminism. But here we are in exactly the dual situation: We allow several equally possible truth-values, and we do not care, which of them is chosen—thus we have *don't care* nondeterminism.

Since our quantifiers depend on the size of the solution set of the quantified constraint, we also have to deal with the situation when this solution set is not measurable. Fortunately, this again is *don't know* nondeterminism. If the solution set is not measurable, then we only know that its volume is between the inner and the outer measure, but we do not know which of these.

A naive approach to modeling such a situation, where a logical formula can have more than one solution set, would propagate sets of truth-values by applying the logical symbols element-wise. For example, if for a sentence $\phi_1 \wedge \phi_2$ the approximate truth-value of ϕ_1 is $\{\mathbf{T}\}$ and the approximate truth-value of ϕ_2 is $\{\mathbf{T}, \mathbf{F}\}$ then the combination of all these elements yields a truth-value of $\{\mathbf{T}, \mathbf{F}\}$ for the whole sentence. This results in a many-valued logic [40, 36]. In order to show that this approach is not feasible, we first demonstrate, that this would need a more complicated many-valued logic, and then give a reason, why to avoid many-valued logics altogether:

One would need a more complicated many-valued logic, because of the

need to consider the interaction between the *don't care* nondeterminism resulting from the approximate quantifiers and the *don't know* nondeterminism resulting from unmeasurable sets. Approximate quantifiers can result in an approximate truth-value $\{\mathbf{T}, \mathbf{F}\}$. An unmeasurable solution set can result in an unknown solution set of a formula. Typically one would model this by an empty set. But one wants to be able to assign the approximate truth-value $\{\mathbf{F}\}$ to the whole formula $\phi_1 \wedge \phi_2$, if ϕ_1 has the approximate truth-value $\{\mathbf{F}\}$ and ϕ_2 has the empty approximate solution set, although element-wise combination yields the empty set in this case. This problem arises because a non-existing truth-value for ϕ_2 means that it can have two possible truth-values ($\mathbf{T}$ or $\mathbf{F}$). We do not know which one, but we have modeled only the *don't care* form of nondeterminism.

One could construct a many-valued logic that solves this problem [17, 15], but there is another problem that makes us avoid such an approach altogether: For example, consider $\phi \wedge \neg\phi$, where ϕ has the approximate solution set $\{\mathbf{T}, \mathbf{F}\}$. Then element-wise combination yields the approximate solution set $\{\mathbf{T}, \mathbf{F}\}$, although we want this formula to be false in any case. The reason is, that a many-valued logic forgets the information about the equality of ϕ in both branches of $\wedge$ [40, 36]. This also makes it impossible to define $\leftrightarrow$ (equivalence) as an abbreviation. A similar problem also occurs in interval mathematics [30, 32], where the information about equal terms is lost.

Before going into the details of our solution, we fix the syntax of the new language. It is the usual one of the first-order predicate language, with the only exception that instead of classical quantifiers, approximate quantifiers are used. Consider the example $\forall_{[0.1, 0.2]}^1 x \exists_{[0.0, 0.1]}^2 y [x > 0 \wedge y = 0]$. The quantifiers have a subscript consisting of a nonnegative real interval (the *annotation*). Furthermore they have a positive integer superscript (the *tag*). We require that, within a formula, quantifiers that have the same tag, also have the same annotation.

Tag equality indicates equal (nondeterministic) behavior of the according quantifiers. That is, for quantifiers that have the same tag, the same element of the corresponding annotations should always be chosen. This will allow us to make $A \wedge \neg A$ false in any case, and makes the definition of $A \leftrightarrow B$ as an abbreviation for $(A \vee \neg B) \wedge (\neg A \vee B)$ possible.

If the annotation of a quantifier is a one-element interval then we say that the quantifier is *deterministic*. A formula where all the tags of quantifiers that are not deterministic, are different, is called *free*. Often we do not explicitly write down the tags but assume an arbitrary tagging such that a formula is free.

For assigning semantics to such formulas, we need to extend some of the usual definitions of the first-order predicate logic:

Definition 5 *An m-structure consists of a measure space $\mathcal{D}$, and for each*

relation and function symbol an according relation and function in $\mathcal{D}$.

As usual, by abuse of notation, we denote by $\mathcal{D}$ also the set on which the measure space is defined. Terms can be interpreted as usual in m-structures. We again fix an arbitrary m-structure $\mathcal{S}$ with measure space $\mathcal{D}$ that defines a measure μ . For defining the semantics of approximate quantification we use a method that is completely orthogonal to the semantics usually given to formulas when we *don't know* the value of certain predicate and function symbols and thus assign validity to the formula using all possible predicate and function symbol assignments.

It is straightforward to give semantics to a first-order constraint with approximate quantifiers if we already know for each occurring quantifier with annotation $[\underline{q}, \bar{q}]$, which volume quantifier $\exists_q$ or $\forall_q$, where $q \in [\underline{q}, \bar{q}]$, and which possible measure to use. So, in analogy to the notion of structure which assigns information to predicate and function symbols, we assign information to quantifiers as follows:

Definition 6 *Given a first-order constraint ϕ , a function $q^* : \mathbb{N} \rightarrow \mathbb{R}$ is a quantifier choice for ϕ iff for every $t \in \mathbb{N}$ that occurs as a quantifier tag in ϕ , $q^*(t) \in [\underline{q}, \bar{q}]$, where $[\underline{q}, \bar{q}]$ is the annotation of the quantifiers occurring in ϕ that are tagged by t .*

In a similar way, we can assign to each tag the possible measure that should be used (recall from Section 2 that in the case of measurable sets a possible measure just assigns its measure, otherwise any value between the inner and the outer measure).

Definition 7 *A measure guess is a function*

$$\mu^* : \mathbb{N} \rightarrow (2^{\mathcal{D}} \rightarrow [0, \infty]),$$

such that for all $t \in \mathbb{N}$, $\mu^(t)$ is a possible measure.*

It is an easy exercise to define the solution set of a constraint ϕ , for a certain quantifier choice q^* for ϕ , and a certain measure-guess μ^* . Now we model that we *don't care* for the quantifier choices and *don't know* about the right measure guesses. For this we introduce two notions corresponding to the notion that a constraint is true. The essence is, that now different quantifier choices can be used, and each choice can result in a different overall result:

Definition 8 *A first-order constraint ϕ is true for a variable assignment θ iff there is a quantifier-choice q^* for ϕ , such that for all measure-guesses μ^* , ϕ is true for θ , q^* , and μ^* . A first-order constraint ϕ is false for a variable assignment θ iff there is a quantifier-choice q^* for ϕ , such that for all measure-guesses μ^* , ϕ is not true for θ , q^* , and μ^* .*

As an example assume a sentence $\exists_{[\underline{q}, \overline{q}]} x \phi$, where ϕ has a measurable solution set d_ϕ . If the volume of d_ϕ is less or equal $\underline{q}$ the constraint is false; if the volume of d_ϕ is greater than $\overline{q}$, then ϕ is true; but if the volume of d_ϕ is within $[\underline{q}, \overline{q}]$, ϕ is both true and false. So we have exactly the behavior sketched in Figure 2.

6 Approximate Computation and Approximate Quantifiers

The question remains, whether we can solve first-order constraints that contain approximate instead of classical quantifiers, up to an arbitrarily small error bound. For this we apply our model for approximate computation of Section 3 to approximate quantification. The agenda will be, first to show how to propagate *all* available approximate information, and then to show, how to infer from this the information needed for our specification in Figure 1.

In Section 3 we represented the uncertainty about a solution set by the notion of approximate solution set of a constraint. In addition to this, here we also have the *don't know* nondeterminism resulting from measure guesses. So an *approximate solution set of a constraint ϕ under a quantifier choice q^** is a potential approximate solution set that contains all solution sets of ϕ under this quantifier choice q^* and any measure guess μ^* . But in addition to this nondeterminism, the semantics of our language, as given in Definition 8, takes into account the *don't care* nondeterminism resulting from quantifier choices. This means that several different approximate solution sets can be equally valid, depending on the actual quantifier choice taken. For modeling this situation we introduce a second level of approximation:

Definition 9 *A potential biapproximate solution set is a set of potential approximate solution sets. A biapproximate solution set of ϕ is a set of approximate solution sets of ϕ .*

Note that here, when dealing with *don't care* nondeterminism, we approximate the case of full information, by taking a subset instead of superset of all possible objects.

In general, also for functions one can have these two forms of nondeterminism: On the one hand, we can have several functions where we *don't know* which is the right one. On the other hand, we can have several functions where we *don't care* which one is chosen. Analogously to solution sets, we model the combination of both by sets of approximate functions, which we call *biapproximate functions*.

For typesetting reasons, instead of $\hat{A}$ we use the notation $\tilde{A}$ to denote biapproximate objects. Again we can deduce the information that we want to find for our specification (Figure 1) as follows:

Definition 10 A variable assignment θ is determined to true by a potential biapproximate solution set $\tilde{d}$ iff there is a potential approximate solution set $\hat{d} \in \tilde{d}$ such that θ is determined to true by $\hat{d}$. It is determined to false by $\tilde{d}$ iff there is a potential approximate solution set $\hat{d} \in \tilde{d}$ such θ is determined to false by $\hat{d}$. It is determined by $\tilde{d}$ iff it is either determined to true or determined to false by $\tilde{d}$.

Now we again define the error of a potential biapproximate solution set $\tilde{d}$ as the outer measure of the set of all variable assignments not determined by $\tilde{d}$.

As for classical quantification also approximate quantification results in a function on potential solution sets:

Definition 11 For a variable $v \in V$, a nonnegative real number q , and a possible measure μ , the (v, q, μ) -projection operator $P_{q, \mu}^v$ is a function on potential solution sets such that $\theta \in P_{q, \mu}^v(d)$ iff $\mu(\{a \in \mathcal{S} \mid \theta_v^a \in d\}) > q$.

For a variable v and nonnegative real number q , we call the set of all (v, q, μ) -projection operators *approximate (v, q) -projection operator* and denote it by P_q^v . This is an approximate function. For a fixed v and interval $[\underline{q}, \overline{q}]$ we call the set of all (v, q) -projection operators, where $q \in [\underline{q}, \overline{q}]$, *biapproximate $(v, [\underline{q}, \overline{q}])$ -projection operator*. This is a biapproximate function.

Now we compute biapproximate solution sets by applying the extension of the functions that correspond to the logical symbols.

Lemma 1 Let $\exists_{[\underline{q}, \overline{q}]}^t v \phi$ be a constraint, $\hat{d}_\phi$ an approximate solution set of ϕ under a quantifier choice q^* for $\exists_{[\underline{q}, \overline{q}]}^t v \phi$, and $P = P_{q^*(t)}^v$. Then $\text{Ext}_P(\hat{d}_\phi)$ is an approximate solution set of $\exists_{[\underline{q}, \overline{q}]}^t v \phi$ under q^* .

Theorem 1 Let $\exists_{[\underline{q}, \overline{q}]}^t v \phi$ be a constraint such that the tag t does not occur in ϕ , let $\tilde{d}_\phi$ be a biapproximate solution set of ϕ , and let P be the biapproximate $(v, [\underline{q}, \overline{q}])$ -projection operator. Then $\text{Ext}_P(\tilde{d}_\phi)$ is a biapproximate solution set of $\exists_{[\underline{q}, \overline{q}]}^t v \phi$.

Proof. Let $\hat{d}$ be an arbitrary but fixed element of $\text{Ext}_P(\tilde{d})$. We have to prove that $\hat{d}$ is an approximate solution set of $\exists_{[\underline{q}, \overline{q}]}^t v \phi$. By definition of extension we know that there is a $\hat{d}_\phi \in \tilde{d}_\phi$ and a $q \in [\underline{q}, \overline{q}]$ such that $\hat{d} = \text{Ext}_{P_q^v}(\hat{d}_\phi)$. By definition of biapproximate solution set, $\hat{d}_\phi$ is an approximate solution set of ϕ under a quantifier choice q_ϕ^* of ϕ . Let q^* be such that it is equal to q_ϕ^* on all tags occurring in ϕ , and let it assign q to t , which is possible since t does not occur in ϕ . Then, by the previous lemma, $\text{Ext}_{P_{q^*(t)}^v}(\hat{d}_\phi) = \hat{d}$ is an approximate solution set of $\exists_{[\underline{q}, \overline{q}]}^t v \phi$ under q^* . ■

The condition on the tags is always fulfilled for free constraints. If it does not hold, the resulting biapproximate solution set might contain elements that are no approximate solution sets of the constraint. The reason for this is, that the extension does not take into account tag equality in a similar way as interval arithmetic does not take into account equality of variables [30, 32].

7 Representing Biapproximate Solution Sets

Before studying, whether biapproximate projection is convergent, we first study biapproximate solution sets in more details. Observe that different potential biapproximate solution sets can contain exactly the same information we need to find for fulfilling our specification in Figure 1. For example, if we have the biapproximate truth value $\{\{\mathbf{T}\}\}$ of a sentence, then we know that the sentence is true. But, if we have the biapproximate truth-value $\{\{\mathbf{T}\}, \{\mathbf{T}, \mathbf{F}\}\}$, then we have exactly the same information. So there is some interaction between the two forms of nondeterminism that potential biapproximate solution sets do not explicitly take into account, and we can divide the potential biapproximate solution sets into equivalence classes such that each equivalence class element contains the same information.

In this section we show, that in many cases it suffices to implement functions on potential biapproximate solution sets (e.g., projection operators) just on the above equivalence classes. Since the set of equivalence classes has a much lower cardinality than the full set of potential biapproximate solution sets, we can then find a representation that is better suited for computer implementation. Furthermore we gain a valuable tool for the subsequent proof that the biapproximate $[g, \bar{q}]$ -projection operators are convergent, and find interesting insight in the interaction between *don't know* and *don't care* nondeterminism.

Before studying the general case of potential biapproximate solution sets, we start with the easier case of potential approximate solution sets. The information we want to extract are the elements determined to true, and the elements determined to false by a potential approximate solution set $\hat{d}$. These are exactly the elements $\bigcap \hat{d}$, and all elements not in $\bigcup \hat{d}$. This gives us the equivalence relation $\hat{d}_1 \sim \hat{d}_2$ iff $(\bigcap \hat{d}_1, \bigcup \hat{d}_1) = (\bigcap \hat{d}_2, \bigcup \hat{d}_2)$.

Now let us also define an order $\leq$ on functions on potential approximate solution sets such that $f_1 \leq f_2$ iff for all d , $f_1(d) \subseteq f_2(d)$. Also here we can form an equivalence relation on approximate functions by defining $\hat{f}_1 \sim \hat{f}_2$ iff $(\min \hat{f}_1, \max \hat{f}_1) = (\min \hat{f}_2, \max \hat{f}_2)$. We call a function f on potential solution sets *monotonic* iff $d_1 \subseteq d_2$ implies $f(d_1) \subseteq f(d_2)$. In a similar way we define an approximate or biapproximate function to be monotonic iff all its members are monotonic.

For computing with the equivalence classes instead of their members, we

need to know whether all members of equivalence classes behave equally for function application. For this we pick a *canonical representative* from each equivalence class and prove that all the other class members behave in the same way as this representative. We can order potential approximate solution sets by the subset relation, and so we can also use interval notation on them: $[\underline{d}, \overline{d}] := \{d \mid \underline{d} \subseteq d \subseteq \overline{d}\}$. Observe that for any potential approximate solution set $\hat{d}$, the interval $r(\hat{d}) := [\bigcap \hat{d}, \bigcup \hat{d}]$ is equivalent to $\hat{d}$, and for any approximate function $\hat{f}$ the interval $r(\hat{f}) := [\min \hat{f}, \max \hat{f}]$ is equivalent to $\hat{f}$. So we take $r(\hat{d})$ and $r(\hat{f})$ as the canonical representatives. Now we have the following congruence property:

Lemma 2 *For a potential approximate solution set $\hat{d}$ with canonical representative $[\underline{d}, \overline{d}]$ and a monotonic approximate function $\hat{f}$ with canonical representative $[\underline{f}, \overline{f}]$, $r(\text{Ext}_{\hat{f}}(\hat{d})) = [\underline{f}(\underline{d}), \overline{f}(\overline{d})]$.*

Proof. By definition, $r(\text{Ext}_{\hat{f}}(\hat{d}))$ is equal to

$$[\bigcap \text{Ext}_{\hat{f}}(\hat{d}), \bigcup \text{Ext}_{\hat{f}}(\hat{d})]$$

Now, by inserting the definition of $\text{Ext}_{\hat{f}}$ and by monotonicity of $\hat{f}$

$$\bigcap \text{Ext}_{\hat{f}}(\hat{d}) = \bigcap \{f(d) \mid f \in \hat{f}, d \in \hat{d}\} = (\min \hat{f})(\bigcap \hat{d}) = \underline{f}(\underline{d})$$

and

$$\bigcup \text{Ext}_{\hat{f}}(\hat{d}) = \bigcup \{f(d) \mid f \in \hat{f}, d \in \hat{d}\} = (\max \hat{f})(\bigcup \hat{d}) = \overline{f}(\overline{d}).$$

■

The (v, q, μ) -projection operators are monotonic. Furthermore the representation of each approximate (v, q) -projection operator is the interval $[P_{q, \underline{\mu}}^v, P_{q, \overline{\mu}}^v]$. So we can compute the approximate projection of a potential approximate solution set by just computing with the interval bounds of the corresponding representations.

Now we study the general case of potential biapproximate solution sets. In a similar way as potential approximate solution sets, different potential biapproximate solution sets can yield the same information. Also here the information we want to extract are the elements determined to true, and the elements determined to false by a potential biapproximate solution set $\tilde{d}$ (see Definition 10). These are the elements $\bigcup \{\bigcap \hat{d} \mid \hat{d} \in \tilde{d}\}$, and all elements not in $\bigcap \{\bigcup \hat{d} \mid \hat{d} \in \tilde{d}\}$. Again, this gives us an equivalence relation $\tilde{d}_1 \sim \tilde{d}_2$ iff $(\bigcup \{\bigcap \hat{d} \mid \hat{d} \in \tilde{d}_1\}, \bigcap \{\bigcup \hat{d} \mid \hat{d} \in \tilde{d}_1\}) = (\bigcup \{\bigcap \hat{d} \mid \hat{d} \in \tilde{d}_2\}, \bigcap \{\bigcup \hat{d} \mid \hat{d} \in \tilde{d}_2\})$. In a similar way we get an equivalence relation on biapproximate functions by $\tilde{f}_1 \sim \tilde{f}_2$ iff $(\max\{\min \hat{f} \mid \hat{f} \in \tilde{f}_1\}, \min\{\max \hat{f} \mid \hat{f} \in \tilde{f}_1\}) = (\max\{\min \hat{f} \mid \hat{f} \in \tilde{f}_2\}, \min\{\max \hat{f} \mid \hat{f} \in \tilde{f}_2\})$.

Again we pick canonical representatives from the equivalence classes. For any elements $\underline{x}$ and $\bar{x}$ in a domain with a partial order $\leq$, let $\llbracket \underline{x}, \bar{x} \rrbracket := \{[y, y \cup \bar{x}] \mid y \in [\underline{x} \cap \bar{x}, \underline{x}]\}$. We call such an object *biinterval* (algebraically speaking the resulting objects form a bilattice [17, 15]). By forming biintervals we stay within an equivalence class:

Lemma 3 *Let A be a set, let $\tilde{a}$ an element of the corresponding set $\tilde{A}$, and let $(\underline{a}, \bar{a}) = (\bigcup\{\bigcap \hat{a} \mid \hat{a} \in \tilde{a}\}, \bigcap\{\bigcup \hat{a} \mid \hat{a} \in \tilde{a}\})$; then $\llbracket \underline{a}, \bar{a} \rrbracket \sim \tilde{a}$.*

Proof.

$$\begin{aligned} \bigcup\{\bigcap \hat{a} \mid \hat{a} \in \llbracket \underline{a}, \bar{a} \rrbracket\} &= \bigcup\{\bigcap [a, a \cup \bar{a}] \mid a \in [\underline{a} \cap \bar{a}, \underline{a}]\} = \\ &= \bigcup\{a \mid a \in [\underline{a} \cap \bar{a}, \underline{a}]\} = \underline{a} = \bigcup\{\bigcap \hat{a} \mid \hat{a} \in \tilde{a}\} \end{aligned}$$

In a similar way $\bigcap\{\bigcup \hat{a} \mid \hat{a} \in \llbracket \underline{a}, \bar{a} \rrbracket\} = \bigcap\{\bigcup \hat{a} \mid \hat{a} \in \tilde{a}\}$. ■

So again, we denote the canonical representative $\llbracket \bigcup\{\bigcap \hat{d} \mid \hat{d} \in \tilde{d}\}, \bigcap\{\bigcup \hat{d} \mid \hat{d} \in \tilde{d}\} \rrbracket$ of $\tilde{d}$ by $r(\tilde{d})$, and the canonical representative $\llbracket \max\{\min \hat{f} \mid \hat{f} \in \tilde{f}\}, \min\{\max \hat{f} \mid \hat{f} \in \tilde{f}\} \rrbracket$ of $\tilde{f}$ by $r(\tilde{f})$. Also here we can compute with the representatives because of the following congruence property:

Theorem 2 *For a potential biapproximate solution set $\tilde{d}$ and a monotonic biapproximate function $\tilde{f}$, $r(\text{Ext}_{\tilde{f}}(\tilde{d})) = \llbracket \underline{f}(\underline{d}), \bar{f}(\bar{d}) \rrbracket$, where $\llbracket \underline{d}, \bar{d} \rrbracket = r(\tilde{d})$, and $\llbracket \underline{f}, \bar{f} \rrbracket = r(\tilde{f})$.*

Proof. By definition, $r(\text{Ext}_{\tilde{f}}(\tilde{d}))$ is equal to

$$\llbracket \bigcup\{\bigcap \hat{d} \mid \hat{d} \in \text{Ext}_{\tilde{f}}(\tilde{d})\}, \bigcap\{\bigcup \hat{d} \mid \hat{d} \in \text{Ext}_{\tilde{f}}(\tilde{d})\} \rrbracket$$

We have:

$$\begin{aligned} \bigcup\{\bigcap \hat{d} \mid \hat{d} \in \text{Ext}_{\tilde{f}}(\tilde{d})\} &= && \text{def. of Ext} \\ \bigcup\{\bigcap \text{Ext}_{\tilde{f}}(\hat{d}) \mid \hat{f} \in \tilde{f}, \hat{d} \in \tilde{d}\} &= && \text{def. of Ext.} \\ \bigcup\{\bigcap \{f(d) \mid f \in \hat{f}, d \in \hat{d}\} \mid \hat{f} \in \tilde{f}, \hat{d} \in \tilde{d}\} &= && \text{monot.} \\ \bigcup\{(\min \hat{f})(\bigcap \hat{d}) \mid \hat{f} \in \tilde{f}, \hat{d} \in \tilde{d}\} &= && \text{monot.} \\ (\max\{\min \hat{f} \mid \hat{f} \in \tilde{f}\})(\bigcup\{\bigcap \hat{d} \mid \hat{d} \in \tilde{d}\}) &= && \text{def. of r} \\ &= && \underline{f}(\underline{d}) \end{aligned}$$

For the dual case an analogous argument holds. ■

For a biapproximate $[q, \bar{q}]$ -projection operator P , $r(P)$ is the biinterval $\llbracket P_{q, \underline{\mu}}, P_{\bar{q}, \bar{\mu}} \rrbracket$. As a consequence of the above theorem, for every potential biapproximate solution set $\tilde{d}$ we can compute the projection of its equivalence class by just working on the bounds $\underline{d}$ and $\bar{d}$ of $r(\tilde{d})$ to get $\llbracket P_{q, \underline{\mu}}(\underline{d}), P_{\bar{q}, \bar{\mu}}(\bar{d}) \rrbracket$.

8 Biapproximate Projection is Convergent

Now we are ready to prove that for approximate quantifiers we can compute approximate solution sets of constraints up to arbitrarily small error bounds. For this we call a potential biapproximate solution set that contains only measurable solution sets *measurable*.

Definition 12 *Let $\tilde{d}$ be a potential biapproximate solution set. For any variable assignment θ the v -error of $\tilde{d}$ over θ is the outer measure of all $a \in S$ for which θ_v^a is not determined by $\tilde{d}$.*

We first prove that for single points in the free variable space we can attain an arbitrarily small error.

Lemma 4 *For every measurable potential biapproximate solution set $\tilde{d}$, for the biapproximate $(v, [\underline{q}, \overline{q}])$ -projection operator P with $\underline{q} < \overline{q}$, for almost all variable assignments θ such that θ is not determined by $r(\text{Ext}_P(\tilde{d}))$, the v -error of $\tilde{d}$ over θ is greater than $\overline{q} - \underline{q}$.*

Proof. Observe that a variable assignment θ is not determined by a potential biapproximate solution set with representation $[[\underline{d}, \overline{d}]]$ iff $\theta \notin \underline{d}$ and $\theta \in \overline{d}$. Let $r(d) = [[\underline{d}, \overline{d}]]$ and let θ be such that it is not determined by $r(\text{Ext}_P([\underline{d}, \overline{d}]))$, which is $[[P_{\underline{q}, \underline{\mu}}^v(\underline{d}), P_{\overline{q}, \overline{\mu}}^v(\overline{d})]]$ by Theorem 2. So, by definition of projection operator, $\underline{\mu}(\{a \mid \theta_v^a \in \underline{d}\}) \leq \underline{q}$ and $\overline{\mu}(\{a \mid \theta_v^a \in \overline{d}\}) > \overline{q}$. Since $\underline{d}$ and $\overline{d}$ are measurable, by Fubini's theorem, the sets $\{a \mid \theta_v^a \in \underline{d}\}$ and $\{a \mid \theta_v^a \in \overline{d}\}$ are measurable for almost all θ . The θ that are not determined by $[[\underline{d}, \overline{d}]]$ are represented by the set $\{a \mid \theta_v^a \notin \underline{d} \wedge \theta_v^a \in \overline{d}\}$. Clearly the volume of this set is greater than $\overline{q} - \underline{q}$. ■

Since we have shown in the last section that for computing our specification (Figure 1), we can apply the canonical representative of projection operators instead of the projection operators themselves, we now prove:

Theorem 3 *For $\underline{q} < \overline{q}$, and the biapproximate $(v, [\underline{q}, \overline{q}])$ -projection operator P , $r \circ \text{Ext}_P$ restricted to measurable potential biapproximate solution sets is convergent.*

Proof. We have to prove that for all $\varepsilon \in \mathbb{R}^+$, there is a $\delta \in \mathbb{R}^+$ such that for every measurable potential biapproximate solution set $\tilde{d}$ such that the error of $\tilde{d}$ is less than δ , the error of $r(\text{Ext}_P(\tilde{d}))$ is less than ε .

Choose $(\overline{q} - \underline{q})\varepsilon$ for δ , and let $\tilde{d}$ be an arbitrary but fixed measurable potential biapproximate solution set. We derive a contradiction from the assumption that the error of $\tilde{d}$ is less than $(\overline{q} - \underline{q})\varepsilon$, but the error of $r(\text{Ext}_P(\tilde{d}))$ is greater or equal ε .

By Lemma 4, for almost every θ that is not determined by $r(\text{Ext}_P(\tilde{d}))$ the v -error of $\tilde{d}$ over θ is greater than $\bar{q} - \underline{q}$. The measure of these θ is greater or equal ε . So, by Fubini's theorem, the error of $\tilde{d}$ is greater than $(\bar{q} - \underline{q})\varepsilon$, which is contradiction to our assumption. ■

Note that, if unmeasurable sets occur, then we still can ensure that the projection operator is convergent, by making the interval $[\underline{q}, \bar{q}]$ big enough. In general, we can make computation faster by increasing the size of the interval $[\underline{q}, \bar{q}]$, that is, by decreasing the required precision.

In a similar way as the projection operators for quantifiers, also for the other logical symbols ($\vee$, $\wedge$, $\neg$) there are corresponding functions on potential solution sets [34], and for free constraints we can use their extension to propagate the according potential biapproximate solution sets. The functions corresponding to conjunction and disjunction are monotonic, and for the function N corresponding to negation, $d_1 \subseteq d_2$ implies $N(d_1 \supseteq d_2)$, which entails a similar property as Theorem 2. So we can use biinterval representation also here.

As an example take the constraint $x^2 + y^2 \leq 1$ and the biinterval $[\emptyset, S_1]$, where the potential solution set S_1 is the set of all variable assignments that assign elements of $[-1, 1]$ to both x and y . This determines all elements outside of the rectangle $[-1, 1] \times [-1, 1]$ to false and leaves the other elements undetermined. Furthermore, take the constraint $x \geq 0$ and the biinterval $[\emptyset, S_2]$, where S_2 is the set of all variable assignments. This leaves all variable assignments undetermined. The solution set of a conjunction is the intersection of the solution sets of the according sub-constraints, and intersection is monotonic. So by Theorem 2 we can take the intersection of the corresponding biinterval elements, which is $\emptyset \cap \emptyset$ and $S_1 \cap S_2$, and get the biinterval $[\emptyset, S_1]$ as a biapproximate solution set for $x^2 + y^2 \leq 1 \wedge x \geq 0$.

One can easily show that the functions corresponding to $\vee$, $\wedge$, and $\neg$ are convergent. So we can compute approximate solution sets of free constraints from approximate solution sets of their atomic sub-constraints, as described in Section 4. Furthermore—as long as no unmeasurable solution sets occur and provided that for all approximate quantifiers the left bound $\underline{q}$ of the annotation is strictly smaller than the right bound $\bar{q}$ —we can attain arbitrarily small, user-defined, error.

9 Related Work

Languages for modeling nondeterminism in various other forms have been introduced within the frame of abstract data types—see for example [43, 42, 21, 26]. There one uses nondeterministic specifications to either model nondeterminism occurring in reality, or to abstract away unnecessary details of the behavior of a real or desired system; these details might be specified

later. Also in analysis nondeterminism has been modeled in order to deal with unknown/uncertain knowledge, and a large amount of classical analysis has been extended to this case [2, 1].

In contrast to the above cases, in our work the deterministic (exact) specification is already given, and we introduce nondeterminism only later, in order to be able to do approximate computation for a relaxed specification. Furthermore we deal with two forms of nondeterminism at the same time whereas the above approaches are always confined to one form of nondeterminism.

The idea to allow several equally valid outputs (i.e., *don't care* nondeterminism) to make certain problems computable is frequently used when doing exact numerical computation on the reals via potentially infinite representations [9, 47, 44].

Modifications of the first-order language that allow reasoning about the size of sets have been studied coming from logic [29], or knowledge representation [3, 4], and they are a main topic in the area of generalized quantifiers [31, 41]. However, these languages do not allow any nondeterministic choice of the size specification and they circumvent the problem of how to deal with unmeasurable sets by allowing only expressions whose solution sets are measurable.

10 Conclusion

For constraints containing classical quantifiers, information about approximate solution sets of the atomic sub-constraints of a first-order constraint does not suffice to compute arbitrarily precise approximate solution sets of the whole constraints. We have provided a remedy for this problem by replacing the classical quantifiers in the first-order predicate language by approximate quantifiers.

In addition to enabling approximation algorithms, this gives us both expressive power in reasoning and the possibility of tunable algorithms (algorithms where the user can decide about the tradeoff between speed and precision). We have implemented such an algorithm—a detailed description and analysis of this implementation will be published elsewhere.

The question remains, how big we should choose the intervals of the quantifier annotation. Too small intervals can hamper the efficiency of constraint solving, while too big intervals can disturb the information one is looking for.

Most of this research was done as a part of the author's Ph.D. work. The author thanks his Ph.D. advisor, Hoon Hong, for all the guidance.

References

- [1] J.-P. Aubin. *Viability Theory*. Birkhäuser, 1991.
- [2] J.-P. Aubin and H. Frankowska. *Set-valued Analysis*. Birkhäuser, Boston, 1990.
- [3] F. Bacchus. Lp, a logic for representing and reasoning with statistical knowledge. *Computational Intelligence*, 6:209–231, 1990.
- [4] F. Bacchus. *Representing and Reasoning with Probabilistic Knowledge*. MIT Press, 1990.
- [5] F. Benhamou, F. Goualard, É. Languénou, and M. Christie. Universally quantified constraint solving: an application to camera control. Technical Report 00.5, Institut de Recherche en Informatique de Nantes, 2000. Submitted for Publication.
- [6] E. Bishop. *Foundations of Constructive Analysis*. McGraw Hill, 1967.
- [7] E. Bishop and D. Bridges. *Constructive Analysis*. Springer, 1985.
- [8] B. Bouchon-Meunier and V. Kreinovich. From interval computations to modal mathematics: Applications and computational complexity. *ACM SIGSAM Bulletin*, 32(2):7–11, 1998.
- [9] L. Brouwer. Besitzt jede reelle Zahl eine Dezimalbruchentwicklung? *Mathematische Annalen*, 83:201–210, 1920.
- [10] B. F. Caviness and J. R. Johnson, editors. *Quantifier Elimination and Cylindrical Algebraic Decomposition*. Springer, Wien, 1998.
- [11] J. H. Davenport and J. Heintz. Real quantifier elimination is doubly exponential. *Journal of Symbolic Computation*, 5:29–35, 1988.
- [12] A. Dolzmann, T. Sturm, and V. Weispfenning. Real quantifier elimination in practice. In B. H. Matzat, G.-M. Greuel, and G. Hiss, editors, *Algorithmic Algebra and Number Theory*, pages 221–248. Springer, 1998.
- [13] P. Dorato, W. Yang, and C. Abdallah. Robust multi-objective feedback design by quantifier elimination. *Journal of Symbolic Computation*, 24:153–159, 1997.
- [14] M. J. Fischer and M. O. Rabin. Super-exponential complexity of presburger arithmetic. *SIAM-AMS Proceedings*, 7:27–41, 1974. Also in [10].
- [15] M. Fitting. Bilattices and the semantics of logic programming. *Journal of Logic Programming*, 11(2):91–116, 1991.
- [16] E. Gardeñes, M. Á. Sainz, L. Jorba, R. Calm, R. Estela, H. Mielgo, and A. Trepát. Modal intervals. *Reliable Computing*, 7(2):77–111, 2001.
- [17] M. L. Ginsberg. Multivalued logics: A uniform approach to reasoning in artificial intelligence. *Computational Intelligence*, 4:265–316, 1988.
- [18] L. Gonzalez-Vega. A combinatorial algorithm solving some quantifier elimination problems. In Caviness and Johnson [10].
- [19] P. Halmos. *Measure Theory*. Van Nostrand, 1950.

- [20] E. Hehner. Predicative programming. *Communications of the ACM*, 27:134–151, Feb. 1984.
- [21] W. H. Hesselink. A mathematical approach to nondeterminism in data types. *ACM Transactions on Programming Languages and Systems*, 10(1):87–117, 1988.
- [22] C. Hoare and H. Jifeng. *Unifying Theories of Programming*. Series in Computer Science. Prentice Hall, 1998.
- [23] H. Hong. Quantifier elimination for formulas constrained by quadratic equations via slope resultants. *The Computer Journal*, 36(5):440–449, 1993. Special issue on computational quantifier elimination.
- [24] H. Hong. Symbolic-numeric methods for quantified constraint solving. In *International Symposium on Scientific Computing, Computer Arithmetic and Validated Numerics SCAN-95*, 1995. Invited Talk.
- [25] H. Hong, R. Liska, and S. Steinberg. Testing stability by quantifier elimination. *Journal of Symbolic Computation*, 24(2):161–187, 1997.
- [26] H. Hussmann. *Nondeterminism in Algebraic Specifications and Algebraic Programs*. Birkhäuser, 1993.
- [27] L. Jaulin, M. Kieffer, O. Didrit, and E. Walter. *Applied Interval Analysis, with Examples in Parameter and State Estimation, Robust Control and Robotics*. Springer, Berlin, 2001.
- [28] M. Jirstrand. *Algebraic Methods for Inequality Constraints in Control*. PhD thesis, Linköping University, Department of Electrical Engineering, Division of Automatic Control, 1998. No. 527.
- [29] H. J. Keisler. Probability quantifiers. In J. Barwise and S. Feferman, editors, *Model Theoretic Logics*, chapter XIV. Springer Verlag, New York, 1985.
- [30] R. E. Moore. *Interval Analysis*. Prentice Hall, Englewood Cliffs, NJ, 1966.
- [31] A. Mostowski. On a generalization of quantifiers. *Fundamenta Mathematicae*, 44:12–36, 1957.
- [32] A. Neumaier. *Interval Methods for Systems of Equations*. Cambridge Univ. Press, Cambridge, 1990.
- [33] M. B. Pour-El and J. I. Richards. *Computability in Analysis and Physics*. Springer-Verlag, 1989.
- [34] S. Ratschan. Uncertainty propagation in heterogeneous algebras for approximate quantified constraint solving. *Journal of Universal Computer Science*, 6(9), 2000.
- [35] S. Ratschan. Applications of quantified constraint solving over the reals—bibliography. <http://www.mpi-sb.mpg.de/~ratschan/appqcs.html>, 2001.
- [36] N. Rescher. *Many-valued Logic*. McGraw-Hill, New York, 1969.
- [37] D. Richardson. Some undecidable problems involving elementary functions of a real variable. *Journal of Symbolic Logic*, 33:514–520, 1968.

- [38] S. P. Shary. Algebraic approach to the interval linear static identification, tolerance, and control problems, or one more application of Kaucher arithmetic. *Reliable Computing*, 2(1):3–33, 1996.
- [39] A. Tarski. *A Decision Method for Elementary Algebra and Geometry*. Univ. of California Press, Berkeley, 1951. Also in [10].
- [40] A. Urquhart. Many-valued logic. In D. Gabbay and F. Guenther, editors, *Handbook of Philosophical Logic, Vol. III: Alternatives in Classical Logic*, chapter III.2, pages 71–116. D. Reidel Publishing Company, 1986.
- [41] J. Väänänen. Generalized quantifiers. *Bulletin of the European Association for Theoretical Computer Science*, June 1997.
- [42] M. Walicki and S. Meldal. A complete calculus for the multialgebraic and functional semantics of nondeterminism. *ACM Transactions on Programming Languages and Systems*, 17(2):366–393, 1995.
- [43] M. Walicki and S. Meldal. Algebraic approaches to nondeterminism: An overview. *ACM Computing Surveys*, 29(1):30–81, 1997.
- [44] K. Weihrauch. *Introduction to Computable Analysis*. Texts in Theoretical Computer Science. Springer, Heidelberg, 2000.
- [45] V. Weispfenning. The complexity of linear problems in fields. *Journal of Symbolic Computation*, 5(1–2):3–27, 1988.
- [46] V. Weispfenning. Quantifier elimination for real algebra — the cubic case. In *Proc. ISSAC’94*, pages 258–263, Oxford, 1994. ACM Press.
- [47] E. Wiedmer. Computing with infinite objects. *Theoretical Computer Science*, 10:133–155, 1980.